

HEAVY QUARKS AND STRUCTURE FUNCTIONS

R. S. THORNE

*Department of Physics, Theoretical Physics, 1 Keble Road,
Oxford OX1 3NP, England
E-mail: thorne@v2.rl.ac.uk*

I present a method for calculating heavy quark production which extrapolates smoothly from the well understood limits of the fixed-flavour number scheme for $Q^2 \sim m_H^2$ to the zero-mass scheme at $Q^2/m_H^2 \rightarrow \infty$. For all $Q^2 > m_H^2$ the evolution of the heavy quark distribution is precisely as in the massless $\overline{\text{MS}}$ scheme. The method is simple to implement and compares well with data.

1 Introduction

Until recently the treatment of heavy flavours in global analyses of structure functions has been rather naive. Usually the charm and bottom quarks were regarded as infinitely massive below $Q^2 = m_H^2$, and massless above this threshold. This procedure, known as the zero mass variable flavour number scheme (ZM-VFNS), gives the correct results for $Q^2/m_H^2 \rightarrow \infty$, but has errors of $\mathcal{O}(m_H^2/Q^2)$ and is very inaccurate near threshold. Alternatively, all heavy flavour final states may be thought of as being produced by the hard scatter between the electroweak boson and a light parton, i.e. the number of active flavours, n_f is 3 and the heavy flavour cross-section is generated (mainly) by photon-gluon fusion. This is called the fixed flavour number scheme (FFNS), and incorporates the correct threshold behaviour, i.e. a smooth kinematic threshold of $W^2 = Q^2(x^{-1} - 1) = 4m_H^2$, automatically. This method is certainly appropriate provided Q^2 is not too large. However, it does not sum potentially large logarithms in Q^2/m_H^2 , and for this reason (and also because of the related lack of a heavy flavour parton distribution) is unsuitable for $Q^2 \gg m_H^2$.

Recently there have been measurements of the charm structure function at HERA^{1,2}, complementing the previous measurements by EMC³. Also, the charm contribution to the total F_2 at small x at HERA can be $> 20\%$, and the NMC data, which contains only 5 – 10% charm, but has very small errors, is surprisingly sensitive to the treatment of charm. Hence, a global analysis of the most up to date structure function data must necessarily include a description of heavy flavours which is satisfactory over a wide range of x and Q^2 .

2 Theoretical Procedure

We desire an approach which extrapolates smoothly from the FFNS at low Q^2 to the ZM-VFNS at high Q^2 , maintaining the correct ordering in both schemes. In order to do this we first examine the relationship between the parton densities in these two schemes (putting $\mu^2 = Q^2$ for simplicity). One can show that the 4 flavour partons in the ZM-VFNS distribution are related to those in the 3 flavour FFNS by the equation

$$f_k^4(Q^2, m_c^2) = A_{ka}(Q^2/m_c^2) \otimes f_a^3(Q^2), \quad (1)$$

where the matrix elements $A_{ka}(Q^2/m_c^2)$ are calculable in perturbation theory, and contain logs in Q^2/m_c^2 .^a Hence, the charm density is totally defined in terms of the light parton densities. Due to the formal equality of the two schemes for $Q^2/m_c^2 \rightarrow \infty$, in this limit we find a reciprocal relationship for the coefficient functions

$$C_{2,ca}^{FF}(Q^2/m_c^2) = C_{2,ck}^4 \otimes A_{ka}(Q^2/m_c^2). \quad (2)$$

So in this limit all the logs in $C_{2,ca}^{FF}(Q^2/m_c^2)$ can be factored into the matrix elements $A_{ka}(Q^2/m_c^2)$, and absorbed into the definition of the 4-flavour parton densities where they are summed automatically by evolution⁴.

For the scales $Q^2 < m_c^2$ we assume the validity of the FFNS approach. For $Q^2 > m_c^2$ we extend the approach outlined above to define a variable flavour number scheme (VFNS). We make precisely the same definition of the 4-flavour parton densities as above (thus ensuring that they evolve as massless partons), but in order to incorporate threshold effects correctly, i.e. guarantee exact equivalence to the FFNS, we have mass dependent coefficient functions. In particular we demand that for all $Q^2 > m_c^2$ our coefficient functions satisfy

$$C_{2,ca}^{FF}(Q^2/m_c^2) = C_{2,ck}^{VF}(Q^2/m_c^2) \otimes A_{ka}(Q^2/m_c^2). \quad (3)$$

Unfortunately, because there are more degrees of freedom in our coefficient functions than there are equations determining them, they are not defined uniquely. For example at lowest order in α_s we have the equation⁵

$$C_{2,cg}^{FF,1}(Q^2/m_c^2) = C_{2,cc}^{VF,0}(Q^2/m_c^2) \otimes A_{cg}(Q^2/m_c^2) + C_{2,cg}^{VF,1}(Q^2/m_c^2). \quad (4)$$

Since in the correct ordering of $F_{2,c}(x, Q^2)$ only the zeroth order coefficient functions appear at LO, the first order coefficient functions appearing at NLO,

^aAt NLO (but not beyond) the charm distribution starts from zero at $Q^2 = m_c^2$ and all other partons are the same in both schemes at this scale.




Figure 1: Comparison of $F_{2,c}(x, Q^2)$ using different schemes.

the way in which we choose to solve (4), and higher order equations, affects our order-by-order expressions for $F_{2,c}(x, Q^2)$. We remove this ambiguity by imposing the physical requirement that at the transition point $Q^2 = m_c^2$, where we go from a 3-flavour to 4-flavour description, $\partial F_{2,c}(x, Q^2)/\partial \ln Q^2$ is continuous. Imposing this to all orders (in the gluon sector) uniquely determines all coefficient functions and defines our VFNS.^b Imposing this constraint, the LO coefficient function is given by the relatively simple expression

$$\partial C_{2,cg}^{FF,1}(Q^2/m_c^2)/\partial \ln Q^2 = C_{2,cc}^{VF,0}(Q^2/m_c^2) \otimes P_{qg}^0, \quad (5)$$

where P_{qg}^0 is the LO quark-gluon splitting function. This then automatically defines $C_{2,cg}^{VF,1}(Q^2/m_c^2)$ via (4). $C_{2,cc}^{VF,1}(Q^2/m_c^2)$ is more complicated, but numerically unimportant. The treatment for bottom quarks is essentially identical, and the procedure can easily be generalized to other processes. Further details can be found in⁶.

Results

Fig. 1 shows the results of the NLO calculation of $F_{2,c}(x, Q^2)$, illustrating how it smoothly extrapolates from the FFNS result at $Q^2 \sim m_c^2$ to the ZM-VFNS

^bWe note that both our coefficient functions and our definition of ordering is different to the alternative, currently LO, VFNS⁵.




Figure 2: Prediction for $F_{2,c}(x, Q^2)$ compared to data.

at high Q^2 . We have also performed global fits using different prescriptions for heavy flavours and find that the best quality fit is indeed obtained using our VFNS approach⁶. Fig. 2 shows the prediction for $F_{2,c}(x, Q^2)$ obtained from the parton distributions, resulting from our best fit to the total $F_2(x, Q^2)$ (using $m_c = 1.35\text{GeV}$), for a variety of values of m_c . Again one can see the smoothness and the good agreement with the full range of available data.

Thus, our VFNS for heavy quark production is on a firm theoretical footing, it is quite simple to implement, clearly has the behaviour we would expect, and matches well to data. Hence we encourage its use in analyses of structure functions.

References

1. H1 collaboration: C. Adloff *et al.*, *Z. Phys. C* **72** 593 (1996).
2. ZEUS collaboration: J. Breitweg *et al.*, *Phys. Lett. B* **407** 402 (1997); Paper N-645 presented at International Europhysics Conference on High Energy Physics, HEP97, Jerusalem 1997.
3. J.J. Aubert *et al.*, *Nucl. Phys. B* **213** 31 (1983).
4. M. Buza *et al.*, *Eur. Phys. J. C* **1** 301 (1998).
5. M. Aivazis *et al.*, *Phys. Rev. D* **50** 3085 (1983).
6. R.S. Thorne and R.G. Roberts, hep-ph/9709442, *Phys. Rev. D* **57** (1998); R.S. Thorne and R.G. Roberts, *Phys. Lett. B* **421** 303 (1998).